

Letters

RESEARCH LETTER


Association Between Observer-Rated Disfigurement and Body Image-Related Distress Among Head and Neck Cancer Survivors

Head and neck cancer (HNC) is associated with a high rate of body image-related distress (BID),¹ a disorder characterized by a distressing self-perceived change in appearance and/or function that can lead to devastating psychosocial morbidity. Prior studies have characterized demographic, clinical, and psychosocial factors associated with HNC-related BID in an attempt to identify high-risk patients and understand the mechanisms underlying this disorder.² However, the contribution of observer-rated disfigurement to BID among HNC survivors remains poorly characterized.³ Therefore, this study's objective is to assess the association between observer-rated disfigurement and BID in HNC survivors.

Methods | This cross-sectional study analyzed data for HNC survivors from 6 institutions from November 2020 to August 2021.¹ The study included 250 patients (see the eTable in the Supplement for demographic and clinical characteristics), was approved by the institutional review board at each participating institution, and followed the STROBE reporting guideline. Written informed consent was obtained from participants. Observer-rated disfigurement was assessed by a clinician using the validated observer-rated disfigurement scale (ORDS; score range, 1-9; higher scores reflecting greater disfigurement).⁴ Interrater reliability for ORDS scores is 0.91 to 0.98.^{3,4} Head and neck cancer-related BID was measured using the Inventory to Measure and Assess imaGe disturbance-Head & Neck (IMAGE-HN), a validated patient-reported outcome measure of HNC-related BID (score range, 0-84; higher scores reflect more severe HNC-related BID).⁵ To characterize the association of ORDS with IMAGE-HN scores, we constructed an unadjusted linear regression model and a multivariable linear regression model adjusted for sex, education, years since treatment completion, and employment (variables associated with IMAGE-HN scores).² Statistical analyses were performed using SAS statistical software, version 9.4 (SAS Institute). Data were analyzed between January 3 and March 14, 2022.

Results | The Figure depicts the ORDS and IMAGE-HN score for each patient. The associations between ORDS and IMAGE-HN scores from the unadjusted and adjusted models are shown in the Table. The unadjusted model shows a positive association between ORDS and IMAGE-HN scores ($\beta = 1.8$; 95% CI, 1.0-2.6). The coefficients of multiple determination (R^2) for the unadjusted model accounted for a minimal proportion of variance (R^2 , 0.08). After adjusting for variables associated with

Figure. Observer-Rated Disfigurement Scale (ORDS) vs Inventory to Measure and Assess imaGe disturbance-Head & Neck (IMAGE-HN) Scores



This scatterplot shows the association between observer-rated disfigurement (measured by ORDS scores) and body image-related distress (measured by IMAGE-HN scores) for 250 survivors of head and neck cancer. Each participant's ORDS value has been jittered horizontally by adding a random number between -0.5 and 0.5 to prevent overlap of data points along the x-axis. For a given level of observer-rated disfigurement (ORDS score), there is significant variability in the corresponding level of body image-related distress (IMAGE-HN score).

Table. Results From Unadjusted and Adjusted Linear Regression Models Estimating IMAGE-HN Scores From ORDS Scores

Variable	β (95% CI)	
	Unadjusted model	Adjusted model
ORDS	1.8 (1.0 to 2.6)	1.6 (0.8 to 2.3)
Post treatment, y	NA	-0.1 (-1.1 to 0.9)
Sex	NA	
Male	NA	1 [Reference]
Female	NA	5.4 (0.6 to 10.1)
Education	NA	
Less than high school	NA	1 [Reference]
Some college	NA	-1.5 (-7.1 to 4.1)
College graduate	NA	-3.9 (-9.8 to 2.0)
Graduate school	NA	-6.6 (-13.3 to 0.4)
Employment	NA	
Employed ^a	NA	1 [Reference]
Not employed ^b	NA	4.4 (-2.0 to 10.7)
Retired	NA	-6.6 (-11.4 to -1.8)
Model R^2	0.08	0.19

Abbreviations: IMAGE-HN, Inventory to Measure and Assess imaGe disturbance-Head & Neck; NA, not applicable; ORDS, observer-rated disfigurement scale.

^a Either part- or full-time or homemaker.

^b Disability or unemployed.

IMAGE-HN scores, there was no association between ORDS and IMAGE-HN scores ($\beta = 1.6$; 95% CI, 0.8-2.3).

Discussion | Using a clinically valid measure of HNC-related BID, we found that observer-rated disfigurement correlated poorly with BID among HNC survivors. This finding adds to prior studies showing minimal association between observer-rated disfigurement and BID among HNC survivors³ as well as other cancer survivor populations (eg, breast cancer).⁵ With its large sample size, multi-institutional nature, and use of an HNC-specific BID patient-reported outcome measure, our study also improves upon methodological limitations of prior studies.³

There is significant heterogeneity in the severity of BID among HNC survivors. As demonstrated in the Figure, for a given level of observer-rated disfigurement, there is a wide range of BID severity. This variability in BID severity, as well as the lack of association with observer-rated disfigurement, is inadequately explained by existing models of HNC-related BID, which have focused on clinical, demographic, and a limited number of psychosocial factors.² Future research should identify factors that moderate the association between disfigurement and BID among HNC survivors.

An important implication of our study is that observer-rated disfigurement should not be used to identify HNC survivors for referral for management of their BID. It may be necessary to develop formal screening protocols and instruments that can be implemented into clinical workflow to accurately identify HNC survivors with BID who may benefit from psychosocial interventions such as cognitive behavioral therapy.⁶ A limitation of this study includes the potential for bias from using 1 observer at each site to rate ORDS, although prior studies indicate excellent interrater reliability.^{3,4} In conclusion, this study shows that among HNC survivors, observer-rated measures of disfigurement correlate poorly with patient-reported BID.

David Macias, MD
 Brittany N. Hand, PhD, OTR/L
 Joseph Zenga, MD
 Patrik Pipkorn, MD, MSCI
 Marci L. Nilsen, PhD, RN
 Amy M. Williams, PhD
 Evan M. Graboyes, MD, MPH

Author Affiliations: Department of Otolaryngology–Head and Neck Surgery, Medical University of South Carolina, Charleston, South Carolina (Macias, Graboyes); School of Health and Rehabilitation Sciences, The Ohio State University, Columbus, Ohio (Hand); Department of Otolaryngology and Communication Sciences, Medical College of Wisconsin, Milwaukee, Wisconsin (Zenga); Department of Otolaryngology–Head and Neck Surgery, Washington University School of Medicine, St Louis, Missouri (Pipkorn); Department of Otolaryngology–Head and Neck Surgery, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania (Nilsen); Department of Family Medicine, Henry Ford Health System, Detroit, Michigan (Williams); Department of Public

Health Sciences, Medical University of South Carolina, Charleston, South Carolina (Graboyes).

Accepted for Publication: March 24, 2022.

Published Online: May 12, 2022. doi:10.1001/jamaoto.2022.0822

Corresponding Author: David Macias, MD, 135 Rutledge Ave, MSC 550, Charleston, SC 29425 (maciasd@musc.edu).

Author Contributions: Drs Macias and Hand had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Macias, Pipkorn, Graboyes.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Macias.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Hand.

Obtained funding: Graboyes.

Administrative, technical, or material support: Macias, Zenga, Williams.

Supervision: Macias, Pipkorn, Williams, Graboyes.

Conflict of Interest Disclosures: Dr Nilsen reported receiving grants from the National Institute of Dental and Craniofacial Research outside the submitted work. Dr Graboyes reported receiving grants from National Cancer Institute during the conduct of the study and grants from the National Cancer Institute and Triologic Society/American College of Surgeons and personal fees from Castle Biosciences and the National Cancer Institute outside the submitted work. No other disclosures were reported.

Funding/Support: This work was supported by R21CA245941 from the National Cancer Institute (Dr Graboyes), DDCF2015209 from the Doris Duke Charitable Foundation (Dr Graboyes), and UL1TR000062 from the National Center for Advancing Translational Sciences.

Role of the Funder/Sponsor: The funding organizations had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: Evan M. Graboyes is a member of the editorial board for *JAMA Otolaryngology–Head & Neck Surgery*, but he was not involved in any of the decisions regarding review of the manuscript or its acceptance.

1. Macias D, Hand BN, Pipkorn P, et al. Association of Inventory to Measure and Assess Image Disturbance–Head and Neck scores with clinically meaningful body image-related distress among head and neck cancer survivors. *Front Psychol*. 2021;12:794038. doi:10.3389/fpsyg.2021.794038

2. Macias D, Hand BN, Maurer S, et al. Factors associated with risk of body image-related distress in patients with head and neck cancer. *JAMA Otolaryngol Head Neck Surg*. 2021;147(12):1019-1026. doi:10.1001/jamaoto.2021.1378

3. Chen SC, Huang CY, Huang BS, et al. Factors associated with healthcare professional's rating of disfigurement and self-perceived body image in female patients with head and neck cancer. *Eur J Cancer Care (Engl)*. 2018;27(2):e12710. doi:10.1111/ecc.12710

4. Katz MR, Irish JC, Devins GM, Rodin GM, Gullane PJ. Reliability and validity of an observer-rated disfigurement scale for head and neck cancer patients. *Head Neck*. 2000;22(2):132-141. doi:10.1002/(SICI)1097-0347(200003)22:2<132::AID-HED4>3.0.CO;2-K

5. Graboyes EM, Hand BN, Ellis MA, et al. Validation of a novel, multidomain head and neck cancer appearance- and function-distress patient-reported outcome measure. *Otolaryngol Head Neck Surg*. 2020;163(5):979-985. doi:10.1177/0194599820927364

6. Graboyes EM, Maurer S, Park Y, et al. Evaluation of a novel telemedicine-based intervention to manage body image disturbance in head and neck cancer survivors. *Psychooncology*. 2020;29(12):1988-1994. doi:10.1002/pon.5399